

An introduction to the volume conjecture

Hitoshi Murakami

Tokyo Institute of Technology

colored Jones polynomial

- $J_N(K; q)$: colored Jones polynomial of a knot K .

colored Jones polynomial

- $J_N(K; q)$: colored Jones polynomial of a knot K .
 - ◆ N : dimension of the irreducible representation of $sl(2, \mathbb{C})$.

colored Jones polynomial

- $J_N(K; q)$: colored Jones polynomial of a knot K .
 - ◆ N : dimension of the irreducible representation of $sl(2, \mathbb{C})$.
 - ◆ q : complex parameter.

colored Jones polynomial

- $J_N(K; q)$: colored Jones polynomial of a knot K .
 - ◆ N : dimension of the irreducible representation of $sl(2, \mathbb{C})$.
 - ◆ q : complex parameter.
 - ◆ $J_2(K; q)$ is the ordinary Jones polynomial.

colored Jones polynomial


- $J_N(K; q)$: colored Jones polynomial of a knot K .
 - ◆ N : dimension of the irreducible representation of $sl(2, \mathbb{C})$.
 - ◆ q : complex parameter.
 - ◆ $J_2(K; q)$ is the ordinary Jones polynomial.
 - ◆ $J_N(\text{unknot}; q) = 1$.

colored Jones polynomial

- $J_N(K; q)$: colored Jones polynomial of a knot K .
 - ◆ N : dimension of the irreducible representation of $sl(2, \mathbb{C})$.
 - ◆ q : complex parameter.
 - ◆ $J_2(K; q)$ is the ordinary Jones polynomial.
 - ◆ $J_N(\text{unknot}; q) = 1$.
 - ◆ a linear combination of $(N - 1)$ -parallel or less of K .

colored Jones polynomial

- $J_N(K; q)$: colored Jones polynomial of a knot K .
 - ◆ N : dimension of the irreducible representation of $sl(2, \mathbb{C})$.
 - ◆ q : complex parameter.
 - ◆ $J_2(K; q)$ is the ordinary Jones polynomial.
 - ◆ $J_N(\text{unknot}; q) = 1$.
 - ◆ a linear combination of $(N - 1)$ -parallel or less of K .

$$\blacksquare J_N(K; q) \approx \sum_{\substack{0 \leq i, j, k, l \leq N-1 \\ i, j, k, l: \text{labelings}}} \left(\prod_{\text{crossings}} R_{kl}^{ij} \right)$$


R-matrix



$$R_{kl}^{ij} := \sum_{m=0}^{\min(N-1-i, j)} \delta_{l, i+m} \delta_{k, j-m} \\ \times (q^{1/2} - q^{-1/2})^m \frac{[i+m]! [N-1+m-j]!}{[i]! [m]! [N-1-j]!} \\ \times q^{(i-(N-1)/2)(j-(N-1)/2) - m(i-j)/2 - m(m+1)/4},$$

R-matrix

$$R_{kl}^{ij} := \sum_{m=0}^{\min(N-1-i, j)} \delta_{l, i+m} \delta_{k, j-m} \\ \times (q^{1/2} - q^{-1/2})^m \frac{[i+m]! [N-1+m-j]!}{[i]! [m]! [N-1-j]!} \\ \times q^{(i-(N-1)/2)(j-(N-1)/2) - m(i-j)/2 - m(m+1)/4},$$

$$[k] := \frac{q^{k/2} - q^{-k/2}}{q^{1/2} - q^{-1/2}}.$$

Volume Conjecture

Volume Conjecture (R. Kashaev, J. Murakami+HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K)$$

Volume Conjecture

Volume Conjecture (R. Kashaev, J. Murakami+HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K)$$

$$S^3 \setminus K \underset{\text{JSJ decomposition}}{=} \left(\bigcup_i H_i \right) \cup \left(\bigcup_j S_j \right).$$

Volume Conjecture

Volume Conjecture (R. Kashaev, J. Murakami+HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K)$$

$$S^3 \setminus K \underset{\text{JSJ decomposition}}{=} \left(\bigcup_i H_i \right) \cup \left(\bigcup_j S_j \right).$$

- H_i : hyperbolic piece,

Volume Conjecture

Volume Conjecture (R. Kashaev, J. Murakami+HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K)$$

$$S^3 \setminus K \underset{\text{JSJ decomposition}}{=} \left(\bigcup_i H_i \right) \cup \left(\bigcup_j S_j \right).$$

- H_i : hyperbolic piece,
- S_j : Seifert fibered piece (S^1 -bundle over a surface with singular fibers).

Volume Conjecture

Volume Conjecture (R. Kashaev, J. Murakami+HM)


$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K)$$

$$S^3 \setminus K \underset{\text{JSJ decomposition}}{=} \left(\bigcup_i H_i \right) \cup \left(\bigcup_j S_j \right).$$



- H_i : hyperbolic piece,
- S_j : Seifert fibered piece (S^1 -bundle over a surface with singular fibers).

$$\text{Vol}(S^3 \setminus K) := \sum_{H_i: \text{hyperbolic pieces}} \text{Vol}(H_i)$$


Jaco–Shalen–Johanson decomposition



Jaco–Shalen–Johanson decomposition




=



Seifert fibered

∪



hyperbolic

Figure-eight knot

- $E :=$ figure-eight knot





Figure-eight knot

- $E :=$ figure-eight knot



(K. Habiro and T. Le)

$$J_N(E; q) = \sum_{k=0}^{N-1} \prod_{j=1}^k \left(q^{(N+j)/2} - q^{-(N+j)/2} \right) \left(q^{(N-j)/2} - q^{-(N-j)/2} \right).$$

Colored Jones polynomial of at the N -th root of unity

$$J_N(E; \exp(2\pi\sqrt{-1}/N)) = \sum_{k=0}^{N-1} \prod_{j=1}^k g_j,$$

Colored Jones polynomial of at the N -th root of unity

$$J_N(E; \exp(2\pi\sqrt{-1}/N)) = \sum_{k=0}^{N-1} \prod_{j=1}^k g_j,$$

where

$$g_j := 4 \sin^2(j\pi/N).$$

Colored Jones polynomial of at the N -th root of unity

$$J_N(E; \exp(2\pi\sqrt{-1}/N)) = \sum_{k=0}^{N-1} \prod_{j=1}^k g_j,$$

where

$$g_j := 4 \sin^2(j\pi/N).$$


We also put

$$f_k := \prod_{j=1}^k g_j$$

so that

$$J_N(E; \exp(2\pi\sqrt{-1}/N)) = \sum_{k=1}^{N-1} f_k$$

Graphs of g_j and $\log f_k$



Proof of the Volume Conjecture for by T. Ekhholm (1)

$$f_{5N/6} < \sum_{k=0}^{N-1} f_k < N f_{5N/6} .$$
$$\parallel$$
$$J_N(E; \exp(2\pi\sqrt{-1}/N))$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (1)

$$f_{5N/6} < \sum_{k=0}^{N-1} f_k < N f_{5N/6} .$$
$$\parallel$$
$$J_N(E; \exp(2\pi\sqrt{-1}/N))$$

\Rightarrow

$$\frac{\log f_{5N/6}}{N} < \frac{\log J_N(E; e^{2\pi\sqrt{-1}/N})}{N} < \frac{\log N}{N} + \frac{\log f_{5N/6}}{N} .$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (1)

$$f_{5N/6} < \sum_{k=0}^{N-1} f_k < N f_{5N/6} .$$
$$\parallel$$
$$J_N(E; \exp(2\pi\sqrt{-1}/N))$$

\Rightarrow

$$\frac{\log f_{5N/6}}{N} < \frac{\log J_N(E; e^{2\pi\sqrt{-1}/N})}{N} < \frac{\log N}{N} + \frac{\log f_{5N/6}}{N} .$$

We use

$$\lim_{N \rightarrow \infty} \frac{\log N}{N} = 0 .$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (2)

$$\Rightarrow \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \lim_{N \rightarrow \infty} \frac{\log f_{5N/6}}{N}$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (2)

$$\begin{aligned} \Rightarrow \quad & \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \lim_{N \rightarrow \infty} \frac{\log f_{5N/6}}{N} \\ & = 2 \lim_{N \rightarrow \infty} \sum_{j=1}^{5N/6} \frac{\log(2 \sin(j\pi/N))}{N} \end{aligned}$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (2)

$$\begin{aligned} \Rightarrow \quad & \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \lim_{N \rightarrow \infty} \frac{\log f_{5N/6}}{N} \\ & = 2 \lim_{N \rightarrow \infty} \sum_{j=1}^{5N/6} \frac{\log(2 \sin(j\pi/N))}{N} \\ & = \frac{2}{\pi} \int_0^{5\pi/6} \log(2 \sin x) dx \end{aligned}$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (2)

$$\Rightarrow \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \lim_{N \rightarrow \infty} \frac{\log f_{5N/6}}{N}$$

$$= 2 \lim_{N \rightarrow \infty} \sum_{j=1}^{5N/6} \frac{\log(2 \sin(j\pi/N))}{N}$$

$$= \frac{2}{\pi} \int_0^{5\pi/6} \log(2 \sin x) dx$$

■ $L(\theta) := - \int_0^\theta \log |2 \sin x| dx$ (Lobachevsky function)

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (2)

$$\Rightarrow \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \lim_{N \rightarrow \infty} \frac{\log f_{5N/6}}{N}$$

$$= 2 \lim_{N \rightarrow \infty} \sum_{j=1}^{5N/6} \frac{\log(2 \sin(j\pi/N))}{N}$$

$$= \frac{2}{\pi} \int_0^{5\pi/6} \log(2 \sin x) dx$$

■ $L(\theta) := - \int_0^\theta \log |2 \sin x| dx$ (Lobachevsky function)

$$= -\frac{2}{\pi} L(5\pi/6)$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (2)

$$\Rightarrow \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \lim_{N \rightarrow \infty} \frac{\log f_{5N/6}}{N}$$


$$= 2 \lim_{N \rightarrow \infty} \sum_{j=1}^{5N/6} \frac{\log(2 \sin(j\pi/N))}{N}$$

$$= \frac{2}{\pi} \int_0^{5\pi/6} \log(2 \sin x) dx$$


■ $L(\theta) := - \int_0^\theta \log |2 \sin x| dx$ (Lobachevsky function)

$$= -\frac{2}{\pi} L(5\pi/6) = \frac{6L(\pi/3)}{2\pi}.$$

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (3)

$$\text{Vol} \left(\begin{array}{c} \text{Diagram} \end{array} \right) = L(\alpha) + L(\beta) + L(\gamma)$$


Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (3)

$$\text{Vol} \left(\text{Diagram} \right) = L(\alpha) + L(\beta) + L(\gamma)$$


$$\Rightarrow \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \frac{6L(\pi/3)}{2\pi}$$

is the volume of the complement of \mathfrak{S} .

Proof of the Volume Conjecture for \mathfrak{S} by T. Ekholm (3)

$$\text{Vol} \left(\text{Diagram} \right) = L(\alpha) + L(\beta) + L(\gamma)$$

$$\Rightarrow \lim_{N \rightarrow \infty} \frac{\log J_N(E; \exp(2\pi\sqrt{-1}/N))}{N} = \frac{6L(\pi/3)}{2\pi}$$

is the volume of the complement of \mathfrak{S} .

- The complement of \mathfrak{S} can be decomposed into two **regular** ideal hyperbolic tetrahedra. So the Volume Conjecture is true for \mathfrak{S} .

The Volume Conjecture is true for ...

The Volume Conjecture is known to be true for

The Volume Conjecture is true for ...

The Volume Conjecture is known to be true for

- Figure-eight knot (T. Ekholm).

The Volume Conjecture is true for ...

The Volume Conjecture is known to be true for

- Figure-eight knot (T. Ekholm).
 - Torus knots (R. Kashaev and O. Tirkkonen).
- Note that Vol of these knots are zero, since the knot complements are Seifert fibered.

The Volume Conjecture is true for ...

The Volume Conjecture is known to be true for

- Figure-eight knot (T. Ekholm).
- Torus knots (R. Kashaev and O. Tirkkonen).
Note that Vol of these knots are zero, since the knot complements are Seifert fibered.
- Whitehead doubles of torus knots of type $(2, p)$ (H. Zheng).

The Volume Conjecture is true for ...

The Volume Conjecture is known to be true for

- Figure-eight knot (T. Ekholm).
- Torus knots (R. Kashaev and O. Tirkkonen).
Note that Vol of these knots are zero, since the knot complements are Seifert fibered.
- Whitehead doubles of torus knots of type $(2, p)$ (H. Zheng).
- Twisted Whitehead links (H. Zheng).

The Volume Conjecture is true for ...

The Volume Conjecture is known to be true for

- Figure-eight knot (T. Ekholm).
- Torus knots (R. Kashaev and O. Tirkkonen).
Note that Vol of these knots are zero, since the knot complements are Seifert fibered.
- Whitehead doubles of torus knots of type $(2, p)$ (H. Zheng).
- Twisted Whitehead links (H. Zheng).
- Borromean rings (S. Garoufalidis and T. Lê).

'Proof' of the Volume Conjecture (1)

$$J_N(K; q) \approx \sum_{\text{labelings}} \prod_{\text{crossings}} R_{kl}^{ij}$$

'Proof' of the Volume Conjecture (1)

$$J_N(K; q) \approx \sum_{\text{labelings}} \prod_{\text{crossings}} R_{kl}^{ij}$$

$\Rightarrow J_N(K; \exp(2\pi\sqrt{-1}/N))$ would be dominated by the
'biggest' $\prod_{\text{crossings}} R_{kl}^{ij}$.

'Proof' of the Volume Conjecture (1)

$$J_N(K; q) \approx \sum_{\text{labelings}} \prod_{\text{crossings}} R_{kl}^{ij}$$

$\Rightarrow J_N(K; \exp(2\pi\sqrt{-1}/N))$ would be dominated by the
'biggest' $\prod_{\text{crossings}} R_{kl}^{ij}$.

F_λ : the biggest one (λ is the labeling).

'Proof' of the Volume Conjecture (1)

$$J_N(K; q) \approx \sum_{\text{labelings}} \prod_{\text{crossings}} R_{kl}^{ij}$$

$\Rightarrow J_N(K; \exp(2\pi\sqrt{-1}/N))$ would be dominated by the 'biggest' $\prod_{\text{crossings}} R_{kl}^{ij}$.

F_λ : the biggest one (λ is the labeling).


We could expect

$$\frac{\log |J(K; \exp(2\pi\sqrt{-1}/N))|}{N} \underset{N \rightarrow \infty}{\sim} \frac{\log F_\lambda}{N}.$$

'Proof' of the VC (2) – Yokota's observation

$$\frac{\log F_\lambda}{N} = \sum_{\text{crossings}} \frac{\log R_{kl}^{ij}}{N}.$$

'Proof' of the VC (2) – Yokota's observation

$$\frac{\log F_\lambda}{N} = \sum_{\text{crossings}} \frac{\log R_{kl}^{ij}}{N}. \quad (\text{biggest one})$$


Y. Yokota observed (after D. Thurston) that for each crossing


'Proof' of the VC (2) – Yokota's observation

$$\frac{\log F_\lambda}{N} = \sum_{\text{crossings}} \frac{\log R_{kl}^{ij}}{N}. \quad (\text{biggest one})$$


Y. Yokota observed (after D. Thurston) that for each crossing

$$\frac{\log R_{kl}^{ij}}{N} \underset{N \rightarrow \infty}{\sim}$$

'Proof' of the VC (2) – Yokota's observation


$$\frac{\log F_\lambda}{N} = \sum_{\text{crossings}} \frac{\log R_{kl}^{ij}}{N}. \quad (\text{biggest one})$$


Y. Yokota observed (after D. Thurston) that for each crossing

$$\frac{\log R_{kl}^{ij}}{N} \underset{N \rightarrow \infty}{\sim}$$

sum of the volumes of the four hyperbolic tetrahedra around the crossing

'Proof' of the VC (2) – Yokota's observation

$$\frac{\log F_\lambda}{N} = \sum_{\text{crossings}} \frac{\log R_{kl}^{ij}}{N}. \quad (\text{biggest one})$$



Y. Yokota observed (after D. Thurston) that for each crossing

$$\frac{\log R_{kl}^{ij}}{N} \underset{N \rightarrow \infty}{\sim}$$

sum of the volumes of the four hyperbolic tetrahedra around the crossing

- The labels would correspond to the complete hyperbolic structure of the knot complement (if the knot has a complete hyperbolic structure).

Four tetrahedra around a vertex



What's wrong?

Why Yokota's method doesn't work even for a hyperbolic knot:

What's wrong?

Why Yokota's method doesn't work even for a hyperbolic knot:

- We don't know whether the method above gives the genuine limit.

What's wrong?

Why Yokota's method doesn't work even for a hyperbolic knot:

- We don't know whether the method above gives the genuine limit.
- To find the 'biggest term', we have to solve a system of equations that corresponds to the hyperbolicity condition of the knot complement.

What's wrong?

Why Yokota's method doesn't work even for a hyperbolic knot:

- We don't know whether the method above gives the genuine limit.
 - To find the 'biggest term', we have to solve a system of equations that corresponds to the hyperbolicity condition of the knot complement.
- ⇒ We have several solutions corresponding to the stationary points.

What's wrong?

Why Yokota's method doesn't work even for a hyperbolic knot:

- We don't know whether the method above gives the genuine limit.
 - To find the 'biggest term', we have to solve a system of equations that corresponds to the hyperbolicity condition of the knot complement.
- ⇒ We have several solutions corresponding to the stationary points.

The problem is which solution gives the 'biggest one'.

Complexification

Complexification (J. Murakami, M. Okamoto,
T. Takata, Y. Yokota, HM)

Complexification

Complexification (J. Murakami, M. Okamoto,
T. Takata, Y. Yokota, HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K).$$

Complexification

Complexification (J. Murakami, M. Okamoto, T. Takata, Y. Yokota, HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log \left| J_N \left(K; \exp(2\pi\sqrt{-1}/N) \right) \right|}{N} = \text{Vol}(S^3 \setminus K).$$

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N \left(K; \exp(2\pi\sqrt{-1}/N) \right)}{N} \\ = \text{Vol}(S^3 \setminus K) + \sqrt{-1} \text{CS}(S^3 \setminus K).$$

Complexification

Complexification (J. Murakami, M. Okamoto, T. Takata, Y. Yokota, HM)

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; \exp(2\pi\sqrt{-1}/N))|}{N} = \text{Vol}(S^3 \setminus K).$$

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N(K; \exp(2\pi\sqrt{-1}/N))}{N} = \text{Vol}(S^3 \setminus K) + \sqrt{-1} \text{CS}(S^3 \setminus K).$$

- One may regard this as a definition of the Chern–Simons invariant for a (non-hyperbolic) knot.

Generalization to three-manifolds

Generalization to three-manifold

Generalization to three-manifolds

Generalization to three-manifold

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N (K; \exp(2\pi\sqrt{-1}/N))}{N} \\ = \text{Vol}(S^3 \setminus K) + \sqrt{-1} \text{CS}(S^3 \setminus K).$$

Generalization to three-manifolds

Generalization to three-manifold

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N (K; \exp(2\pi\sqrt{-1}/N))}{N} \\ = \text{Vol}(S^3 \setminus K) + \sqrt{-1} \text{CS}(S^3 \setminus K).$$

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N (K; \exp(a/N))}{N} = ?$$

Generalization to three-manifolds

Generalization to three-manifold

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N (K; \exp(2\pi\sqrt{-1}/N))}{N} = \text{Vol}(S^3 \setminus K) + \sqrt{-1} \text{CS}(S^3 \setminus K).$$

$$2\pi \lim_{N \rightarrow \infty} \frac{\log J_N (K; \exp(a/N))}{N} = ?$$

to be continued...